

Ship To: In Office Infusion Suite At Home Other _____

PATIENT INFORMATION

Patient Name:		SSN:		DOB:	
Address:		City:		State:	Zip:
Home Phone:		Height:	Weight:	Gender:	Male Female
Cell Phone:		Email Address:			

INSURANCE INFORMATION (or attach copy of cards)

Primary Insurance Co:	Policy Holder:	Relationship:	Policy #:	Group #:
Secondary Insurance:	Policy Holder:	Relationship:	Policy #:	Group #:

CLINICAL INFORMATION (Fax all pertinent clinical and lab information)

Diagnosis (ICD-10): _____ **Date of Diagnosis:** _____

Pemphigus L10.0	CIDP G61.81	Peripheral Neuropathy G60.9	MMN G61.82	Multiple Sclerosis G35	Acute Infective Polyneuritis/GBS G61.0
Myasthenia Gravis with acute exacerbation G70.01		Myasthenia Gravis without acute exacerbation G70.00		Dermatomyositis M33.90	Polymyositis M33.20

Has patient received immune globulin previously? No Yes: Date of last infusion _____ Date of next infusion: _____

Comorbidities: _____

ALLERGIES: NKDA Other _____

PRESCRIPTION INFORMATION (or attach a copy of the prescription)

Infusion Therapy:
 Preferred brand _____ **OR** Pharmacist will determine appropriate product based on clinical assessment, insurance requirements and availability)
 No Substitute Refills: _____ times (as allowed by state or payer requirements)

Dose: (please select option(s) and provide complete information, pharmacy to round the nearest 5 gram vial)
 Administration Rate = Follow Manufacturer's Guidelines
 Loading Dose: _____ gm/kg over _____ days, then
 Maintenance dose: _____ gm/kg over _____ days, every _____ weeks x _____ cycles
 Other Regimen _____

Infusion Rate: (please select one and provide complete information)
 Pharmacist to determine
 Start at _____ ml/hr, then increase by _____ ml/hr every _____ minutes to maximum

Vascular Access Device:
 Peripheral Catheter PICC Port
 Other (describe/# of lumens): _____

Flush Orders: (If IV ordered the following flush protocols will be followed)
 Sodium Chloride 0.9%
 Peripheral Line: 3 ml before each dose and 3 ml after each dose and prn
 Central Line: 5 - 10 ml before each dose and 5 -10 ml after each dose and prn
 Heparin 10 u/ml Peripheral Line: 3 ml after last sodium flush and prn
 Heparin 100 u/ml Central Line: 5 ml after last sodium flush and prn
 Provide needles, syringes, VAD supplies & other ancillary supplies needed for infusion

Pre-Medication
Diphenhydramine
 25 mg capsule: 1-2 capsules by mouth 15-30 minutes before each infusion Decline

Acetaminophen
 650 mg tablet: 1-2 tablets by mouth 15-30 minutes before each infusion Decline
 Other _____ Strength: _____
 Directions: _____

Hydration Orders
 Infuse _____ mg _____ solution
 Prior to Following

Labs:
 Results will be faxed to physician's office. If no frequency noted, ordered labs to be done prior to initial infusion only. Labs will not be drawn on weekend/holidays. Not appropriate for STAT labs
 Quantitative Ig A prior to first dispense. Pharmacist to obtain authorization from MD.
 Other _____ Frequency of Labs: _____

As required by your state, Prescriber to check "Dispense as written" or handwritten "Brand Medically Necessary" and sign to prevent generic substitution. Dispense as written

NURSING

Nursing Agency: _____ **Phone:** _____
 Skilled Nursing Visits for Immune Globulin Intravenous administration and education. To provide education related to the disease process and IG therapy. To provide an assessment of patient's general overall health status. To provide skilled nursing visits PRN for additional education and support.

PHYSICIAN INFORMATION

Prescriber Name:	Phone:	Fax:
Office Contact:	Email:	
Address:		
NPI #:	Tax ID #	
Prescriber Signature:	Date	

725765

Sales Person: